

**Relating Respiratory Disease mortality
to previous smoking habits.
Within-country analysis**

Country Index

1	Australia
2	Austria
3	Belgium
4	Bulgaria
5	Canada
6	Czechoslovakia
7	Denmark
8	Finland
9	France
10	Germany
11	Greece
12	Hungary
13	Iceland
14	Ireland
15	Israel
16	Italy
17	Japan
18	Netherlands
19	New Zealand
20	Norway
21	Poland
22	Portugal
23	Romania
24	Spain
25	Sweden
26	Switzerland
27	UK
28	USA
29	USSR
30	Yugoslavia

Notes for Tables 1 to 30:

Prevalence = Prevalence of smoking (%), total cigarettes, 20 years earlier.

Consumption = Consumption of total cigarettes per adult, 20 years earlier.

Cons tar adj = Consumption of total cigarettes per adult, 20 years earlier, square root tar adjusted.

N = Number of data pairs

R = Pearson correlation coefficient

Summary of results within 30 countries

31	Respiratory Disease mortality and prevalence of smoking, 20 years earlier
32	Respiratory Disease mortality and consumption, 20 years earlier
33	Respiratory Disease mortality and consumption (square root tar adjusted), 20 years earlier
34	Respiratory Disease mortality and prevalence of smoking, averaged from 25 to 5 years earlier
35	Respiratory Disease mortality and consumption, averaged from 25 to 5 years earlier
36	Respiratory Disease mortality and consumption (square root tar adjusted), averaged from 25 to 5 years earlier
37	Respiratory Disease mortality and prevalence of smoking, 20 years earlier – all tobacco products

Table R5.1
Respiratory Diseases mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Australia

	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Males Age 45-49													
Mortality rate	11.6	9.0	9.5	10.9	15.5	15.3	12.4	9.6	8.0	5.5	3.9		
Prevalence										48.2	45.4	2	
Consumption										13.6	12.4	2	
Cons tar adj										9.4	7.9	2	
Males Age 60-64													
Mortality rate	84.7	79.4	103.5	138.5	162.9	151.5	129.6	113.6	105.2	85.9	66.6		
Prevalence										46.3	42.1	2	
Consumption										14.4	13.8	2	
Cons tar adj										10.0	8.8	2	
Males Age 75-79													
Mortality rate	306.0	306.5	415.9	519.1	718.3	830.7	796.1	756.8	685.2	605.6	493.2		
Prevalence										39.2	36.6	2	
Consumption										12.1	10.7	2	
Cons tar adj										8.4	6.8	2	
Females Age 45-49													
Mortality rate	5.6	3.9	4.1	3.6	8.2	11.5	9.1	8.6	7.8	5.6	4.4		
Prevalence										35.7	34.8	2	
Consumption										8.5	8.2	2	
Cons tar adj										5.9	5.2	2	
Females Age 60-64													
Mortality rate	12.1	10.6	15.6	16.8	27.2	38.2	42.9	48.4	54.1	49.6	42.2		
Prevalence										33.6	30.8	2	
Consumption										8.4	7.9	2	
Cons tar adj										5.8	5.0	2	
Females Age 75-79													
Mortality rate	127.0	91.2	82.6	67.2	93.7	104.1	128.4	158.6	211.1	235.8	224.4		
Prevalence										24.9	23.7	2	
Consumption										6.9	6.2	2	
Cons tar adj										4.8	3.9	2	

Table R5.2
Respiratory diseases mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Austria

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		9.3	9.3	7.3	11.3	9.9	8.9	10.2	6.3	5.3	5.9		
Prevalence						93.7	82.4	71.1	59.8	48.5	51.0	6	0.87
Consumption										13.8	14.0	2	
Cons tar adj										10.1	9.1	2	
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		57.7	77.0	82.6	123.6	67.2	54.5	69.1	59.2	51.7	45.9		
Prevalence						85.2	74.8	64.4	54.0	43.6	39.1	6	0.71
Consumption										14.5	14.7	2	
Cons tar adj										10.6	9.6	2	
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		229.5	225.0	352.1	562.9	407.8	392.9	441.2	356.6	324.2	281.5		
Prevalence						75.9	66.7	57.5	48.3	39.1	38.4	6	0.79
Consumption										11.8	12.0	2	
Cons tar adj										8.6	7.8	2	
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		1.7	3.3	3.2	4.8	3.1	3.9	4.4	3.9	3.1	2.7		
Prevalence						40.3	35.2	30.2	25.1	20.0	27.5	6	0.10
Consumption										3.7	5.3	2	
Cons tar adj										2.7	3.5	2	
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		17.8	15.2	17.1	26.4	19.0	19.7	19.3	16.0	14.4	14.0		
Prevalence						28.5	24.5	20.6	16.6	12.6	17.1	6	0.82
Consumption										2.9	4.2	2	
Cons tar adj										2.1	2.7	2	
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		152.4	110.8	160.2	195.2	127.8	103.9	131.4	105.5	108.6	93.8		
Prevalence						17.5	15.2	12.9	10.5	8.2	11.6	6	0.44
Consumption										1.7	2.4	2	
Cons tar adj										1.2	1.6	2	

Table R5.3
Respiratory Diseases mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Belgium

Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		48.7	41.9	33.1	32.2	23.2	18.6	11.7	9.9	10.2			
Prevalence								77.7	74.6	71.6	56.2	3	0.76
Consumption											14.1	0	
Cons tar adj											10.2	0	
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		244.1	225.0	228.2	240.4	181.2	163.3	154.9	145.9	121.5			
Prevalence								70.6	67.8	65.1	51.1	3	0.96
Consumption											14.8	0	
Cons tar adj											10.7	0	
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		405.0	390.4	478.3	643.8	610.7	641.2	717.9	742.7	770.1			
Prevalence								62.9	60.5	58.0	38.9	3	-1.00
Consumption											9.2	0	
Cons tar adj											6.7	0	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		6.0	3.7	5.4	6.1	5.8	6.4	6.9	5.5	5.3			
Prevalence								18.4	15.8	13.1	33.1	3	0.90
Consumption											5.6	0	
Cons tar adj											4.1	0	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		24.0	21.7	22.7	30.7	27.8	24.4	27.2	30.2	35.8			
Prevalence								17.7	13.5	9.3	23.4	3	-0.98
Consumption											4.4	0	
Cons tar adj											3.2	0	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		197.7	150.3	161.9	174.7	154.3	142.3	127.9	143.9	174.1			
Prevalence								13.9	9.8	5.7	14.3	3	-0.98
Consumption											2.0	0	
Cons tar adj											1.5	0	

Table R5.5
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Canada

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	8.5	7.7	8.0	7.9	9.9	11.5	9.8	7.0	5.9	4.9	4.1		
Prevalence							82.6	62.0	58.6	54.5	49.4	5	0.99
Consumption								16.9	16.1	15.3	14.6	4	1.00
Cons tar adj								14.7	12.5	10.9	9.6	4	1.00
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	34.7	40.9	58.1	73.6	94.5	101.3	93.2	76.0	69.8	60.1	50.1		
Prevalence							80.0	65.0	58.7	49.6	44.9	5	0.99
Consumption								17.8	17.0	16.3	15.3	4	1.00
Cons tar adj								15.5	13.2	11.6	10.0	4	0.98
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	125.0	107.2	169.7	217.2	318.3	431.2	452.1	512.3	550.3	508.2	466.9		
Prevalence							75.3	58.6	53.6	48.2	42.9	5	-0.33
Consumption								15.3	14.5	14.5	13.3	4	0.63
Cons tar adj								13.3	11.2	10.3	8.7	4	0.58
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	2.0	3.5	2.9	4.6	5.5	6.1	6.1	5.1	4.0	3.4	3.1		
Prevalence									42.2	41.6	39.6	3	0.91
Consumption								9.0	9.4	9.9	10.4	4	-0.96
Cons tar adj								7.8	7.3	7.0	6.8	4	1.00
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	13.4	10.2	12.9	13.8	19.8	28.1	28.5	31.5	35.9	35.1	34.9		
Prevalence									40.4	39.9	37.9	3	0.76
Consumption								9.0	9.5	9.8	10.4	4	0.63
Cons tar adj								7.8	7.4	7.0	6.8	4	-0.73
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	69.7	49.7	53.1	53.4	66.9	82.0	97.8	130.2	175.6	203.4	217.7		
Prevalence									28.4	29.6	30.0	3	1.00
Consumption								5.9	6.6	7.3	7.9	4	0.98
Cons tar adj								5.1	5.1	5.2	5.2	4	0.74

Table R5.7
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Denmark

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		6.7	3.2	9.0	8.6	13.4	7.9	11.3	5.3	4.1	5.0		
Prevalence						31.2	53.9	49.7	49.9	52.7	51.6	6	-0.75
Consumption						8.5	8.1	7.8	8.6	9.0	9.1	6	-0.66
Cons tar adj						8.5	7.8	7.0	7.8	8.3	7.9	6	-0.10
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		24.6	29.5	70.3	105.1	107.8	104.5	111.4	112.4	95.4	88.9		
Prevalence						25.3	49.1	45.2	45.4	50.2	47.2	6	-0.35
Consumption						7.6	7.9	8.2	8.6	8.9	8.9	6	-0.60
Cons tar adj						7.6	7.6	7.3	7.8	8.2	7.8	6	-0.57
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		74.1	105.6	199.3	271.5	363.2	411.8	519.5	564.6	581.4	661.3		
Prevalence						18.3	43.7	40.3	39.4	40.4	42.2	6	0.62
Consumption						3.2	4.9	6.6	6.4	6.5	7.1	6	0.93
Cons tar adj						3.2	4.7	5.9	5.8	6.0	6.2	6	0.91
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		2.3	2.2	2.5	4.0	7.6	8.4	8.2	7.8	3.9	5.9		
Prevalence						51.1	51.5	42.3	44.6	48.0	49.7	6	-0.14
Consumption						7.4	6.6	5.8	6.6	7.2	7.3	6	-0.58
Cons tar adj						7.4	6.3	5.2	6.0	6.6	6.4	6	-0.31
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		9.4	7.6	14.6	21.6	31.7	42.5	57.0	82.9	108.6	110.4		
Prevalence						32.3	39.9	40.5	42.4	48.0	48.4	6	0.94
Consumption						4.8	5.3	5.8	6.4	7.1	7.1	6	0.99
Cons tar adj						4.8	5.1	5.2	5.8	6.5	6.2	6	0.98
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		51.4	61.0	95.1	91.6	104.2	110.9	129.6	192.3	266.0	365.3		
Prevalence						18.1	26.8	31.8	32.4	33.8	36.3	6	0.76
Consumption						2.4	3.5	4.5	4.6	4.9	5.1	6	0.76
Cons tar adj						2.4	3.4	4.0	4.2	4.5	4.4	6	0.74

Table R5.8
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Finland

Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		10.6	14.4	13.5	14.5	13.7	9.1	5.2	4.4	4.2	2.8		
Prevalence								58.8	55.6	49.8	41.6	4	0.97
Consumption								13.6	12.6	11.6	10.5	4	0.96
Cons tar adj								12.5	10.9	9.2	7.6	4	0.95
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		73.8	115.6	129.3	144.0	118.6	108.7	87.4	75.6	51.9	51.5		
Prevalence							63.9	61.7	56.3	47.9	29.5	5	0.84
Consumption							13.5	14.3	13.5	12.6	8.4	5	0.66
Cons tar adj							13.1	13.2	11.7	10.0	6.1	5	0.83
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		196.1	412.7	456.5	608.6	586.4	594.2	503.4	393.5	368.0	422.3		
Prevalence							61.0	54.4	49.6	39.6	26.4	5	0.69
Consumption							11.0	10.9	9.8	8.6	7.0	5	0.67
Cons tar adj							10.7	10.0	8.5	6.8	5.1	5	0.75
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		1.1	2.5	1.9	3.6	3.7	3.0	3.8	3.0	2.2	2.5		
Prevalence									23.8	30.3	25.6	3	-0.95
Consumption										4.8	4.4	2	
Cons tar adj										3.8	3.2	2	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		7.2	9.7	9.9	15.3	18.2	18.6	21.5	18.1	16.6	18.6		
Prevalence									16.5	17.8	15.8	3	-1.00
Consumption										2.7	3.1	2	
Cons tar adj										2.1	2.2	2	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		30.2	45.9	76.9	98.3	68.7	73.0	72.3	64.8	74.2	94.7		
Prevalence									8.7	11.1	8.5	3	-0.27
Consumption										1.3	1.5	2	
Cons tar adj										1.0	1.1	2	

Table R5.9
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

France

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	24.8	24.0	19.1	15.6	18.3	18.1	15.7	11.6	7.7	6.6	6.4		
Prevalence						75.1	92.0	71.3	67.1	62.9	58.6	6	0.79
Consumption								13.3	12.9	12.4	11.9	4	0.86
Cons tar adj								11.7	10.9	9.7	8.5	4	0.83
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	91.1	94.6	94.2	87.0	98.4	92.0	86.4	76.7	61.6	49.2	46.1		
Prevalence						72.8	83.7	64.8	61.3	57.7	54.2	6	0.88
Consumption								13.9	13.4	12.8	12.3	4	0.97
Cons tar adj								12.2	11.3	10.1	8.8	4	0.94
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	253.0	299.9	330.8	357.0	425.0	473.8	494.5	461.9	348.8	318.2	305.8		
Prevalence						73.2	77.3	57.8	54.0	50.2	46.5	6	0.90
Consumption								11.3	10.5	9.6	8.8	4	0.90
Cons tar adj								9.9	8.9	7.5	6.3	4	0.88
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	7.7	6.9	5.0	4.4	5.9	6.0	5.4	4.9	3.4	3.0	3.3		
Prevalence						23.5	22.1	20.8	26.7	32.7	38.6	6	-0.78
Consumption								1.6	2.3	3.0	3.8	4	-0.77
Cons tar adj								1.4	1.9	2.4	2.7	4	-0.85
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	35.2	32.4	26.3	22.0	23.4	22.6	21.6	20.3	16.8	14.9	15.6		
Prevalence						20.0	17.8	15.5	19.4	23.3	27.2	6	-0.69
Consumption								1.6	2.0	2.5	2.9	4	-0.87
Cons tar adj								1.4	1.7	2.0	2.1	4	-0.94
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	167.3	180.0	170.1	173.8	178.0	167.7	155.4	140.2	110.2	107.6	108.1		
Prevalence						11.0	11.8	12.6	13.9	15.2	16.6	6	-0.92
Consumption								1.7	1.7	1.7	1.6	4	0.35
Cons tar adj								1.5	1.4	1.3	1.1	4	0.63

Table R5.10
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Germany

Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		17.7	13.4	9.6	11.3	12.2	10.7	9.5	6.8	7.4	5.8		
Prevalence					73.5	78.8	74.7	68.8	69.6	58.6	56.2	7	0.88
Consumption											13.2	1	
Cons tar adj											8.3	1	
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		103.3	137.6	142.8	158.3	124.0	104.1	91.1	84.7	85.0	60.1		
Prevalence					66.9	71.7	67.9	62.5	63.3	50.7	47.5	7	0.73
Consumption											13.1	1	
Cons tar adj											8.2	1	
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		331.8	391.5	393.1	569.2	677.6	650.1	596.8	517.4	497.9	414.7		
Prevalence					59.6	63.9	60.5	55.7	56.4	46.1	42.9	7	0.91
Consumption											10.6	1	
Cons tar adj											6.6	1	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		3.2	2.9	3.2	5.2	6.0	6.7	6.1	4.5	4.1	4.0		
Prevalence					31.4	32.0	34.0	30.5	42.5	43.9	43.6	7	-0.85
Consumption											7.6	1	
Cons tar adj											4.8	1	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		20.2	20.4	18.0	25.2	24.5	25.6	25.5	24.7	27.3	22.1		
Prevalence					22.2	22.6	24.0	21.6	30.0	29.7	27.1	7	0.04
Consumption											5.1	1	
Cons tar adj											3.2	1	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		186.0	183.6	141.0	153.3	147.2	127.9	119.6	121.9	141.2	129.3		
Prevalence					13.6	13.9	14.7	13.2	18.5	18.4	18.6	7	-0.26
Consumption											3.2	1	
Cons tar adj											2.0	1	

Table R5.16
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Italy

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		20.9	18.4	18.0	17.6	15.3	11.8	8.4	5.9	4.2	4.3		
Prevalence					82.0	79.3	76.7	52.3	58.3	64.4	69.4	7	0.73
Consumption					10.6	10.1	9.7	9.6	10.8	11.9	11.6	7	-0.58
Cons tar adj					10.6	10.1	9.4	8.9	9.5	9.9	9.0	7	0.70
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		78.0	121.5	155.0	165.6	130.6	115.9	95.4	73.4	53.9	39.5		
Prevalence					79.3	71.4	63.5	56.5	57.5	58.5	65.9	7	0.71
Consumption					11.1	10.8	10.5	10.1	11.4	12.6	13.1	7	-0.70
Cons tar adj					11.1	10.8	10.2	9.4	10.0	10.4	10.1	7	0.55
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		289.4	346.8	440.7	543.7	648.3	633.5	635.0	581.7	498.5	408.7		
Prevalence					63.0	61.4	59.7	51.8	51.9	52.1	60.6	7	-0.10
Consumption					9.0	8.6	8.2	8.2	9.2	10.2	11.1	7	-0.96
Cons tar adj					9.0	8.6	8.0	7.6	8.1	8.4	8.6	7	-0.48
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		5.4	5.2	4.7	4.9	4.7	3.9	3.4	2.8	2.4	2.3		
Prevalence					11.3	9.4	7.4	24.4	24.8	25.3	34.2	7	-0.88
Consumption					0.9	0.7	0.5	0.5	4.2	7.8	4.0	7	-0.78
Cons tar adj					0.9	0.7	0.5	0.5	3.7	6.5	3.1	7	-0.76
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		33.0	34.7	36.2	37.5	29.1	26.6	22.1	18.0	14.4	12.8		
Prevalence					10.8	8.9	7.1	24.0	21.0	17.9	21.6	7	-0.71
Consumption					0.9	0.7	0.5	0.5	3.3	6.1	2.4	7	-0.67
Cons tar adj					0.9	0.7	0.5	0.5	2.9	5.1	1.9	7	-0.64
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		231.8	240.8	251.9	235.9	209.6	176.2	163.4	148.3	131.4	110.7		
Prevalence					8.5	7.0	5.5	17.6	14.3	11.0	13.8	7	-0.56
Consumption					0.7	0.6	0.4	0.4	2.0	3.6	1.5	7	-0.59
Cons tar adj					0.7	0.6	0.4	0.4	1.8	3.0	1.2	7	-0.55

Table R5.17
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Japan

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	19.2	10.1	6.2	5.0	6.7	6.1	5.2	4.9	4.2	4.1	4.0		
Prevalence					81.8	80.8	79.9	78.5	80.1	83.1	81.4	7	-0.02
Consumption								17.4	15.7	18.6	19.9	4	-0.29
Cons tar adj								15.2	12.4	13.6	13.7	4	0.81
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	96.3	55.7	33.9	28.5	47.1	44.6	36.5	36.1	34.6	36.5	32.7		
Prevalence					85.7	82.7	79.7	82.5	81.8	78.5	73.8	7	0.76
Consumption							17.5	18.3	16.6	19.4	20.9	5	-0.47
Cons tar adj							16.2	16.0	13.1	14.2	14.4	5	0.47
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	388.1	292.3	229.3	200.6	322.1	351.2	317.0	332.4	327.3	347.6	347.8		
Prevalence					81.8	78.8	75.7	78.0	77.0	73.0	73.0	7	-0.43
Consumption							15.8	14.9	13.7	16.7	17.0	5	0.62
Cons tar adj							14.6	13.0	10.8	12.2	11.7	5	-0.53
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	9.8	5.2	2.8	2.2	4.2	3.8	3.0	2.9	2.8	2.6	2.5		
Prevalence					10.9	10.9	10.8	8.9	10.0	10.8	16.1	7	-0.28
Consumption								1.2	1.3	1.5	2.1	4	-0.94
Cons tar adj								1.1	1.0	1.1	1.4	4	-0.86
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	46.6	26.1	13.3	10.8	18.7	19.5	15.5	15.0	13.9	13.0	11.8		
Prevalence					13.7	13.7	13.7	12.4	15.5	13.4	14.4	7	-0.24
Consumption								1.7	1.9	2.1	2.6	4	-0.98
Cons tar adj								1.5	1.5	1.5	1.8	4	-0.87
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	199.3	150.6	107.9	91.6	133.0	132.5	108.9	108.9	103.0	107.5	107.5		
Prevalence					17.4	17.4	17.4	16.4	20.0	17.4	16.0	7	-0.15
Consumption								2.3	2.6	2.7	3.2	4	-0.03
Cons tar adj								2.0	2.1	2.0	2.2	4	-0.04

Table R5.19
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

New Zealand

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	14.1	10.1	10.8	12.8	17.5	14.7	18.2	13.9	8.6	6.1	5.9		
Prevalence								55.5	52.5	49.5	42.5	4	0.81
Consumption								17.4	15.3	13.1	11.0	4	0.92
Cons tar adj								16.1	13.4	10.8	8.5	4	0.93
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	56.1	105.7	111.3	151.6	145.8	138.9	152.9	131.1	111.1	85.6	68.9		
Prevalence								58.1	51.5	45.0	43.8	4	0.98
Consumption								18.3	16.3	14.4	12.4	4	1.00
Cons tar adj								16.9	14.3	11.9	9.5	4	1.00
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	187.1	439.8	624.1	711.2	799.1	833.3	911.3	824.1	745.7	644.6	530.8		
Prevalence								55.5	47.8	40.1	56.7	4	0.00
Consumption								14.9	15.1	15.2	15.4	4	-0.99
Cons tar adj								13.8	13.2	12.5	11.8	4	1.00
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	5.6	6.5	6.9	10.7	15.3	15.2	18.4	15.2	14.5	6.6	4.8		
Prevalence								56.7	47.0	37.3	37.8	4	0.91
Consumption								9.9	7.9	5.9	8.1	4	0.61
Cons tar adj								9.2	6.9	4.9	6.2	4	0.78
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	2.5	17.1	21.5	35.2	47.6	52.3	60.6	69.5	77.8	64.3	63.0		
Prevalence								54.3	45.1	35.9	35.3	4	0.57
Consumption								9.9	7.9	5.9	7.9	4	0.31
Cons tar adj								9.2	6.9	4.9	6.1	4	0.43
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	90.9	116.8	156.1	166.1	144.9	157.8	194.4	179.9	232.9	260.3	315.8		
Prevalence								42.6	33.3	24.0	30.6	4	-0.71
Consumption								7.7	6.1	4.6	5.9	4	-0.65
Cons tar adj								7.1	5.3	3.8	4.5	4	-0.80

Table R5.20
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Norway

Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		5.2	4.9	2.8	3.6	5.7	6.2	8.0	5.1	4.7	3.8		
Prevalence						62.8	57.4	52.1	52.0	52.0	54.2	6	0.02
Consumption										8.6	8.7	2	
Cons tar adj										7.3	6.9	2	
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		19.1	27.5	33.6	39.7	48.6	59.0	66.6	53.3	52.7	57.1		
Prevalence						57.1	52.2	47.3	48.2	49.0	49.9	6	-0.60
Consumption										8.8	8.6	2	
Cons tar adj										7.5	6.8	2	
Males	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		69.7	67.2	107.5	136.7	207.1	272.8	313.7	329.5	347.4	390.3		
Prevalence						50.9	46.8	42.8	43.2	43.7	43.2	6	-0.90
Consumption								7.2	7.6	7.9	7.4	4	0.14
Cons tar adj								6.7	6.8	6.7	5.8	4	-0.92
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 45-49													
Mortality rate		2.7	2.0	1.5	2.8	4.7	3.7	5.2	2.9	4.1	4.3		
Prevalence						37.1	39.0	41.0	49.9	58.8	52.4	6	-0.34
Consumption										5.4	5.8	2	
Cons tar adj										4.6	4.6	2	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 60-64													
Mortality rate		9.8	10.7	7.7	12.3	17.3	18.7	25.9	33.6	39.9	42.8		
Prevalence						26.2	27.6	29.0	35.2	41.5	40.7	6	0.98
Consumption										4.5	5.1	2	
Cons tar adj										3.8	4.0	2	
Females	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Age 75-79													
Mortality rate		57.7	65.8	57.7	52.1	72.6	79.3	95.8	101.3	147.1	169.1		
Prevalence						16.1	15.2	14.4	20.0	25.5	28.2	6	0.95
Consumption										3.0	3.2	2	
Cons tar adj										2.5	2.5	2	

Table R5.25
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Sweden

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		2.4	2.3	2.7	4.8	7.2	7.0	5.0	4.7	3.7	3.0		
Prevalence								51.7	46.3	46.5	35.6	4	0.89
Consumption								7.9	4.9	6.7	6.6	4	0.01
Cons tar adj								7.4	4.2	5.3	4.7	4	0.48
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		15.3	14.9	24.5	26.6	38.8	44.4	42.5	35.3	36.8	32.7		
Prevalence								46.2	42.1	40.1	35.0	4	0.91
Consumption								6.8	5.1	5.2	6.5	4	0.35
Cons tar adj								6.4	4.4	4.1	4.6	4	0.82
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		57.6	59.8	99.0	141.2	206.2	251.9	227.0	232.9	238.5	246.0		
Prevalence								39.2	36.2	37.1	29.4	4	-0.90
Consumption								4.2	3.3	4.1	5.0	4	0.62
Cons tar adj								4.0	2.8	3.3	3.5	4	-0.22
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		2.4	1.7	1.2	2.4	4.3	4.6	5.3	3.2	3.2	2.1		
Prevalence								38.8	46.5	48.6	43.3	4	-0.59
Consumption								4.4	4.4	6.4	6.9	4	-0.74
Cons tar adj								4.1	3.8	5.1	4.8	4	-0.45
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		8.4	7.8	8.1	11.6	17.5	23.9	25.9	25.7	30.5	28.9		
Prevalence								29.0	32.9	34.4	33.0	4	0.70
Consumption								3.5	3.5	4.0	5.3	4	0.58
Cons tar adj								3.3	3.0	3.2	3.7	4	0.32
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		33.6	36.1	51.7	63.6	71.5	78.6	71.9	91.0	114.8	132.5		
Prevalence								14.8	19.4	21.9	23.0	4	0.96
Consumption								1.4	1.7	2.6	3.5	4	0.98
Cons tar adj								1.3	1.5	2.1	2.5	4	0.98

Table R5.26
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

Switzerland

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		8.5	10.0	9.9	12.3	10.1	7.1	6.6	3.6	3.5	2.6		
Prevalence							46.3	46.1	45.9	45.7	39.3	5	0.64
Consumption							5.1	9.0	12.9	16.8	13.7	5	-0.89
Cons tar adj							4.9	8.2	11.1	12.1	8.5	5	-0.70
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		38.4	50.7	68.0	84.6	90.7	86.5	75.4	65.2	52.1	42.4		
Prevalence							35.9	37.6	39.3	41.1	34.6	5	-0.07
Consumption							5.4	8.8	12.2	15.6	13.1	5	-0.89
Cons tar adj							5.2	8.1	10.5	11.2	8.1	5	-0.60
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		183.5	186.6	222.6	308.4	355.6	360.8	347.1	349.6	360.2	393.5		
Prevalence							17.7	24.7	31.6	38.6	30.7	5	0.14
Consumption							4.4	7.3	10.2	13.1	8.9	5	0.02
Cons tar adj							4.3	6.7	8.7	9.4	5.5	5	-0.41
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		2.3	2.7	2.2	3.4	4.0	4.6	2.9	2.8	1.8	0.8		
Prevalence										47.7	38.2	2	
Consumption										9.0	9.1	2	
Cons tar adj										6.5	5.6	2	
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		12.7	13.1	11.1	13.7	17.3	16.9	15.9	17.2	19.0	20.5		
Prevalence										27.2	25.5	2	
Consumption										6.4	5.4	2	
Cons tar adj										4.6	3.3	2	
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate		97.3	80.0	78.2	82.1	92.0	70.6	71.3	80.8	92.7	115.6		
Prevalence										19.0	14.2	2	
Consumption										4.3	2.2	2	
Cons tar adj										3.1	1.4	2	

Table R5.27
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

United Kingdom (Total cigarettes)

	1946-1950	1951-1955	1956-1960	1961-1965	1966-1970	1971-1975	1976-1980	1981-1985	1986-1990	1991-1995	1996-2000	N	R
Males													
Age 45-49													
Mortality rate	39.8	38.1	29.5	27.7	23.4	19.0	14.3	10.8	8.4	6.8	6.1		
Prevalence								68.9	62.8	56.6	48.3	4	0.95
Consumption										13.8	11.8	2	
Cons tar adj										10.5	8.3	2	
Males													
Age 60-64													
Mortality rate	288.6	300.1	307.3	317.4	269.5	194.3	154.8	128.1	117.3	93.2	82.4		
Prevalence								62.7	58.7	54.7	47.4	4	0.96
Consumption										14.4	12.7	2	
Cons tar adj										11.0	8.9	2	
Males													
Age 75-79													
Mortality rate	672.6	744.8	800.9	987.1	1013.8	964.3	876.6	791.7	744.8	662.9	596.8		
Prevalence								55.8	54.1	52.3	47.0	4	0.96
Consumption										12.1	11.3	2	
Cons tar adj										9.2	7.9	2	
Females													
Age 45-49													
Mortality rate	9.4	10.0	9.1	10.4	12.3	12.9	10.5	8.1	6.5	5.4	4.9		
Prevalence	18.1	29.0	39.8	41.5	43.2	44.8	46.5	49.0	47.8	49.5	46.6	11	-0.27
Consumption										9.1	8.9	2	
Cons tar adj										6.9	6.2	2	
Females													
Age 60-64													
Mortality rate	60.1	57.2	51.8	51.2	49.0	50.0	52.2	58.4	68.6	59.9	56.1		
Prevalence	17.3	27.8	38.2	41.1	44.1	47.0	49.9	52.9	55.0	53.1	47.2	11	0.09
Consumption										9.2	9.2	2	
Cons tar adj										7.0	6.5	2	
Females													
Age 75-79													
Mortality rate	351.7	316.6	240.4	246.1	208.9	178.5	159.5	168.4	209.9	258.8	300.9		
Prevalence	13.6	21.8	29.9	31.1	32.4	33.7	34.9	39.6	44.9	44.5	43.7	11	-0.52
Consumption										7.4	7.9	2	
Cons tar adj										5.6	5.5	2	

Table R5.27a
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

United Kingdom (Manufactured cigarettes)

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-		
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	N	R
Mortality rate	39.8	38.1	29.5	27.7	23.4	19.0	14.3	10.8	8.4	6.8	6.1		
Prevalence					72.0	68.3	66.2	58.7	59.1	52.1	46.8	7	0.93
Consumption					10.1	11.7	11.3	10.8	11.7	11.8	10.4	7	-0.30
Cons tar adj					9.7	11.0	10.6	9.8	9.9	9.0	7.3	7	0.59
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-		
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	N	R
Mortality rate	288.6	300.1	307.3	317.4	269.5	194.3	154.8	128.1	117.3	93.2	82.4		
Prevalence					71.2	65.6	65.5	59.4	57.4	50.8	43.3	7	0.89
Consumption					11.6	12.8	13.7	12.6	12.4	11.5	10.4	7	0.26
Cons tar adj					11.1	12.0	12.9	11.5	10.5	8.7	7.3	7	0.59
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-		
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	N	R
Mortality rate	672.6	744.8	800.9	987.1	1013.8	964.3	876.6	791.7	744.8	662.9	596.8		
Prevalence					58.8	58.4	58.4	53.0	51.1	46.6	42.4	7	0.96
Consumption					9.5	10.4	11.2	10.3	9.9	9.8	8.8	7	0.47
Cons tar adj					9.1	9.8	10.5	9.4	8.4	7.5	6.2	7	0.80
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-		
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	N	R
Mortality rate	9.4	10.0	9.1	10.4	12.3	12.9	10.5	8.1	6.5	5.4	4.9		
Prevalence					53.1	50.4	52.4	49.6	49.8	49.3	45.8	7	0.77
Consumption					4.2	4.9	5.7	6.1	7.1	8.5	8.1	7	-0.96
Cons tar adj					4.0	4.6	5.4	5.6	6.0	6.5	5.7	7	-0.88
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-		
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	N	R
Mortality rate	60.1	57.2	51.8	51.2	49.0	50.0	52.2	58.4	68.6	59.9	56.1		
Prevalence					43.5	43.0	49.8	53.6	55.0	53.5	46.8	7	0.86
Consumption					3.5	4.3	5.8	6.8	7.8	8.9	7.9	7	0.78
Cons tar adj					3.3	4.0	5.5	6.2	6.6	6.8	5.5	7	0.84
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-		
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000	N	R
Mortality rate	351.7	316.6	240.4	246.1	208.9	178.5	159.5	168.4	209.9	258.8	300.9		
Prevalence					34.1	33.7	39.1	42.0	43.1	42.8	42.9	7	0.44
Consumption					2.7	3.3	4.5	5.2	5.8	6.4	7.4	7	0.69
Cons tar adj					2.6	3.1	4.2	4.7	4.9	4.9	5.2	7	0.44

Table R5.28
Respiratory Disease mortality rates (per 100,000 per year) and smoking habits
20 years earlier, by age, period and sex, showing Pearson correlation coefficients

USA

Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	7.9	8.2	9.7	11.3	13.8	12.9	11.2	9.8	9.3	8.6	8.2		
Prevalence		66.1	60.5	54.9	65.4	62.7	61.7	61.2	56.3	48.6	43.4	10	0.52
Consumption					17.1	17.5	15.8	17.3	16.7	14.8	14.6	7	0.70
Cons tar adj					17.1	17.7	14.9	14.4	13.0	11.0	9.9	7	0.95
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	35.8	44.9	68.2	89.7	116.1	113.9	104.0	100.8	100.5	96.0	90.4		
Prevalence		48.1	50.6	53.1	61.2	59.2	56.2	58.1	52.8	47.1	45.2	10	0.66
Consumption					16.1	18.7	16.5	18.1	18.0	17.2	16.2	7	0.17
Cons tar adj					16.1	19.0	15.6	15.1	14.0	12.8	11.0	7	0.90
Males	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	66.3	84.4	136.9	220.7	337.1	414.5	442.2	516.3	534.7	512.2	495.0		
Prevalence		38.3	40.2	42.2	47.9	45.2	47.1	50.9	45.8	38.3	38.5	10	0.40
Consumption					12.1	12.7	12.5	14.8	14.6	14.0	13.5	7	0.91
Cons tar adj					12.1	12.9	11.8	12.4	11.4	10.4	9.1	7	-0.44
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 45-49	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	2.2	2.5	3.2	4.7	7.2	8.0	7.1	7.7	7.5	7.4	7.4		
Prevalence		26.8	37.2	47.6	43.2	35.9	41.2	41.3	41.1	37.4	35.3	10	0.37
Consumption					8.3	7.2	7.7	9.8	10.5	9.3	9.7	7	-0.02
Cons tar adj					8.3	7.3	7.3	8.2	8.2	6.9	6.6	7	0.06
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 60-64	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	7.1	7.2	9.8	13.7	23.9	31.6	40.4	50.2	60.3	66.9	69.0		
Prevalence		14.8	29.5	44.3	34.8	32.2	40.4	40.0	38.6	37.2	38.3	10	0.51
Consumption					5.6	6.5	7.9	10.3	10.7	11.3	9.3	7	0.89
Cons tar adj					5.6	6.6	7.5	8.6	8.3	8.4	6.3	7	0.53
Females	1946-	1951-	1956-	1961-	1966-	1971-	1976-	1981-	1986-	1991-	1996-	N	R
Age 75-79	1950	1955	1960	1965	1970	1975	1980	1985	1990	1995	2000		
Mortality rate	34.9	32.0	35.9	43.7	54.6	73.3	101.3	162.3	226.4	273.6	304.8		
Prevalence		7.6	15.2	22.8	17.4	13.3	22.0	25.5	25.5	29.1	31.0	10	0.85
Consumption					2.4	2.2	3.9	5.4	6.4	8.5	8.4	7	0.99
Cons tar adj					2.4	2.2	3.7	4.5	5.0	6.3	5.7	7	0.95

Table R5.31
Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to prevalence of smoking 20 years earlier, within 30 countries.

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	2		2		2		2		2		2	
Austria	6	0.87	6	0.71	6	0.79	6	0.10	6	0.82	6	0.44
Belgium	3	0.76	3	0.96	3	-1.00	3	0.90	3	-0.98	3	-0.98
Bulgaria	0		0		0		0		0		0	
Canada	5	0.99	5	0.99	5	-0.33	3	0.91	3	0.76	3	1.00
Czechoslovakia	1		1		2		1		1		1	
Denmark	6	-0.75	6	-0.35	6	0.62	6	-0.14	6	0.94	6	0.76
Finland	4	0.97	5	0.84	5	0.69	3	-0.95	3	-1.00	3	-0.27
France	6	0.79	6	0.88	6	0.90	6	-0.78	6	-0.69	6	-0.92
Germany	7	0.88	7	0.73	7	0.91	7	-0.85	7	0.04	7	-0.26
Greece	5	-0.41	5	0.56	5	0.46	3	-0.69	3	-0.86	3	-0.86
Hungary	3	-0.83	4	0.03	4	0.53	3	0.24	3	0.88	3	-0.83
Iceland	0		0		0		0		0		0	
Ireland	3	0.89	3	0.98	3	0.99	3	0.49	3	0.95	3	-0.86
Israel	4	-0.18	4	0.85	4	0.85	3	0.23	3	0.20	3	-0.99
Italy	7	0.73	7	0.71	7	-0.10	7	-0.88	7	-0.71	7	-0.56
Japan	7	-0.02	7	0.76	7	-0.43	7	-0.28	7	-0.24	7	-0.15
Netherlands	5	0.76	5	0.94	5	-0.35	5	-0.19	5	0.49	5	0.66
NewZealand	4	0.81	4	0.98	4	0.00	4	0.91	4	0.57	4	-0.71
Norway	6	0.02	6	-0.60	6	-0.90	6	-0.34	6	0.98	6	0.95
Poland	5	0.29	5	0.20	5	0.65	5	-0.90	5	-0.81	5	-0.85
Portugal	2		2		2		2		2		2	
Romania	3	-0.88	3	-1.00	3	0.28	3	-0.46	3	-0.74	3	-0.94
Spain	3	-0.97	3	-0.99	3	0.96	3	-0.82	3	-0.98	3	-0.98
Sweden	4	0.89	4	0.91	4	-0.90	4	-0.59	4	0.70	4	0.96
Switzerland	5	0.64	5	-0.07	5	0.14	2		2		2	
UK	4	0.95	4	0.96	4	0.96	11	-0.27	11	0.09	11	-0.52
USA	10	0.52	10	0.66	10	0.40	10	0.37	10	0.51	10	0.85
USSR	0		0		0		0		0		0	
Yugoslavia	2		3	0.20	3	-0.70	1		1		1	
UK (Manufactured Cigs)	7	0.93	7	0.89	7	0.96	7	0.77	7	0.86	7	0.44

Table R5.32
Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to consumption 20 years earlier, within 30 countries.

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	2		2		2		2		2		2	
Austria	2		2		2		2		2		2	
Belgium	0		0		0		0		0		0	
Bulgaria	0		0		0		0		0		0	
Canada	4	1.00	4	1.00	4	0.63	4	-0.96	4	0.63	4	0.98
Czechoslovakia	0		0		0		0		0		0	
Denmark	6	-0.66	6	-0.60	6	0.93	6	-0.58	6	0.99	6	0.76
Finland	4	0.96	5	0.66	5	0.67	2		2		2	
France	4	0.86	4	0.97	4	0.90	4	-0.77	4	-0.87	4	0.35
Germany	1		1		1		1		1		1	
Greece	5	0.35	5	-0.90	5	-0.89	1		1		1	
Hungary	3	-0.82	3	-0.77	3	-0.75	3	0.10	3	0.69	3	-0.96
Iceland	0		0		0		0		0		0	
Ireland	4	-0.09	4	-0.12	4	-0.23	4	-0.16	4	-0.63	4	0.98
Israel	4	0.63	4	0.34	4	0.33	3	-0.45	3	0.27	3	-0.99
Italy	7	-0.58	7	-0.70	7	-0.96	7	-0.78	7	-0.67	7	-0.59
Japan	4	-0.29	5	-0.47	5	0.62	4	-0.94	4	-0.98	4	-0.03
Netherlands	5	0.21	5	-0.61	5	0.23	5	-0.17	5	0.86	5	0.86
NewZealand	4	0.92	4	1.00	4	-0.99	4	0.61	4	0.31	4	-0.65
Norway	2		2		4	0.14	2		2		2	
Poland	5	-0.98	5	-0.95	5	-0.99	5	-0.86	5	-0.86	5	-0.99
Portugal	0		0		0		0		0		0	
Romania	1		1		1		1		1		1	
Spain	3	-0.99	3	0.98	3	-0.97	1		1		1	
Sweden	4	0.01	4	0.35	4	0.62	4	-0.74	4	0.58	4	0.98
Switzerland	5	-0.89	5	-0.89	5	0.02	2		2		2	
UK	2		2		2		2		2		2	
USA	7	0.70	7	0.17	7	0.91	7	-0.02	7	0.89	7	0.99
USSR	0		0		0		0		0		0	
Yugoslavia	0		3	0.00	3	-0.67	0		0		1	
UK (Manufactured Cigs)	7	-0.30	7	0.26	7	0.47	7	-0.96	7	0.78	7	0.69

Table R5.33

Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to consumption (square root tar adjusted) 20 years earlier, within 30 countries.

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	2		2		2		2		2		2	
Austria	2		2		2		2		2		2	
Belgium	0		0		0		0		0		0	
Bulgaria	0		0		0		0		0		0	
Canada	4	1.00	4	0.98	4	0.58	4	1.00	4	-0.73	4	0.74
Czechoslovakia	0		0		0		0		0		0	
Denmark	6	-0.10	6	-0.57	6	0.91	6	-0.31	6	0.98	6	0.74
Finland	4	0.95	5	0.83	5	0.75	2		2		2	
France	4	0.83	4	0.94	4	0.88	4	-0.85	4	-0.94	4	0.63
Germany	1		1		1		1		1		1	
Greece	0		0		0		0		0		0	
Hungary	0		0		0		0		0		0	
Iceland	0		0		0		0		0		0	
Ireland	0		0		0		0		0		0	
Israel	0		0		0		0		0		0	
Italy	7	0.70	7	0.55	7	-0.48	7	-0.76	7	-0.64	7	-0.55
Japan	4	0.81	5	0.47	5	-0.53	4	-0.86	4	-0.87	4	-0.04
Netherlands	0		0		0		0		0		0	
NewZealand	4	0.93	4	1.00	4	1.00	4	0.78	4	0.43	4	-0.80
Norway	2		2		4	-0.92	2		2		2	
Poland	0		0		0		0		0		0	
Portugal	0		0		0		0		0		0	
Romania	0		0		0		0		0		0	
Spain	0		0		0		0		0		0	
Sweden	4	0.48	4	0.82	4	-0.22	4	-0.45	4	0.32	4	0.98
Switzerland	5	-0.70	5	-0.60	5	-0.41	2		2		2	
UK	2		2		2		2		2		2	
USA	7	0.95	7	0.90	7	-0.44	7	0.06	7	0.53	7	0.95
USSR	0		0		0		0		0		0	
Yugoslavia	0		0		0		0		0		0	
UK (Manufactured Cigs)	7	0.59	7	0.59	7	0.80	7	-0.88	7	0.84	7	0.44

Table R5.34

Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to prevalence of smoking averaged from 25 to 5 years earlier, within 30 countries.

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	1		1		1		1		1		1	
Austria	5	0.82	5	0.55	5	0.84	5	-0.27	5	0.87	5	-0.21
Belgium	1		1		1		1		1		1	
Bulgaria	0		0		0		0		0		0	
Canada	4	1.00	4	0.99	4	0.56	2		2		2	
Czechoslovakia	0		0		1		0		0		0	
Denmark	5	0.37	5	0.04	5	0.95	5	0.32	5	0.99	5	0.97
Finland	2		2		3	0.95	1		1		1	
France	5	0.95	5	1.00	5	0.98	5	-0.92	5	-0.79	5	-0.95
Germany	6	0.97	6	0.93	6	0.99	6	-0.86	6	0.44	6	-0.12
Greece	4	0.46	4	0.89	4	0.83	2		2		2	
Hungary	2		2		3	1.00	2		2		2	
Iceland	0		0		0		0		0		0	
Ireland	3	0.92	2		2		2		2		2	
Israel	3	-0.42	3	0.97	3	0.57	2		2		2	
Italy	6	0.99	6	0.98	6	0.93	6	-0.95	6	-0.98	6	-0.87
Japan	6	0.96	6	0.66	6	-0.30	6	-0.51	6	0.87	6	0.36
Netherlands	1		1		2		1		1		1	
NewZealand	3	0.89	3	1.00	3	0.94	3	0.97	3	0.94	3	-0.89
Norway	5	0.70	5	0.33	5	0.15	5	-0.03	5	1.00	5	0.98
Poland	4	0.81	4	0.84	4	0.98	4	-0.97	4	-0.94	4	-0.96
Portugal	1		1		1		1		1		1	
Romania	2		2		2		2		2		2	
Spain	2		2		2		2		2		2	
Sweden	3	1.00	3	0.66	3	-1.00	3	0.98	3	0.80	3	1.00
Switzerland	3	0.79	3	-1.00	3	0.55	0		0		0	
UK	3	0.97	3	0.97	3	0.99	10	0.13	10	0.33	10	-0.24
USA	9	0.78	9	0.20	9	-0.11	9	-0.06	9	0.58	9	0.96
USSR	0		0		0		0		0		0	
Yugoslavia	1		1		2		0		0		0	
UK (Manufactured Cigs)	5	0.98	5	0.95	5	0.96	5	0.99	5	0.70	5	0.67

Table R5.35
Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to consumption averaged from 25 to 5 years earlier, within 30 countries.

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	1		1		1		1		1		1	
Austria	1		1		1		1		1		1	
Belgium	0		0		0		0		0		0	
Bulgaria	0		0		0		0		0		0	
Canada	3	0.99	3	1.00	3	1.00	3	0.60	3	0.63	3	0.99
Czechoslovakia	0		0		0		0		0		0	
Denmark	5	-0.55	5	-0.20	5	0.99	5	-0.66	5	1.00	5	0.95
Finland	2		2		3	0.94	0		0		0	
France	3	0.92	3	0.95	3	0.99	3	-0.25	3	-0.61	3	-0.70
Germany	0		0		0		0		0		0	
Greece	3	1.00	3	-0.97	3	-0.91	0		0		0	
Hungary	2		2		2		2		2		2	
Iceland	0		0		0		0		0		0	
Ireland	3	-0.09	2		2		2		2		2	
Israel	1		1		1		0		0		0	
Italy	6	-0.89	6	-0.74	6	-0.73	6	-0.98	6	-0.99	6	-0.96
Japan	3	-0.98	3	-0.70	4	0.55	3	-0.98	3	-0.90	3	-0.82
Netherlands	4	0.93	4	0.94	4	0.30	4	0.58	4	0.99	4	0.99
NewZealand	1		1		1		1		1		1	
Norway	1		1		3	-0.99	1		1		1	
Poland	3	-0.97	3	-0.96	3	-0.99	3	-0.98	3	-0.96	3	-0.98
Portugal	0		0		0		0		0		0	
Romania	0		0		0		0		0		0	
Spain	2		2		2		0		0		0	
Sweden	3	-0.90	3	-0.86	3	0.98	3	-0.90	3	0.63	3	0.99
Switzerland	2		2		2		0		0		0	
UK	1		1		1		1		1		1	
USA	5	0.85	5	0.65	5	0.44	5	-0.08	5	0.91	5	0.99
USSR	0		0		0		0		0		0	
Yugoslavia	0		0		2		0		0		0	
UK (Manufactured Cigs)	5	0.92	5	0.76	5	0.66	5	-0.96	5	0.93	5	0.73

Table R5.36
Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to consumption (square root tar adjusted) averaged from 25 to 5 years earlier, within 30 countries.

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	0		0		0		0		0		0	
Austria	0		0		0		0		0		0	
Belgium	0		0		0		0		0		0	
Bulgaria	0		0		0		0		0		0	
Canada	1		1		1		1		1		1	
Czechoslovakia	0		0		0		0		0		0	
Denmark	4	0.61	4	0.09	4	0.98	4	0.40	4	1.00	4	0.94
Finland	2		2		3	0.94	0		0		0	
France	3	0.92	3	0.95	3	0.98	3	-0.30	3	-0.63	3	-0.72
Germany	0		0		0		0		0		0	
Greece	0		0		0		0		0		0	
Hungary	0		0		0		0		0		0	
Iceland	0		0		0		0		0		0	
Ireland	0		0		0		0		0		0	
Israel	0		0		0		0		0		0	
Italy	6	0.82	6	0.93	6	0.98	6	-0.99	6	-0.99	6	-0.95
Japan	3	0.93	3	0.36	4	-0.83	3	-0.99	3	0.96	3	-0.86
Netherlands	0		0		0		0		0		0	
NewZealand	1		1		1		1		1		1	
Norway	0		0		0		0		0		0	
Poland	0		0		0		0		0		0	
Portugal	0		0		0		0		0		0	
Romania	0		0		0		0		0		0	
Spain	0		0		0		0		0		0	
Sweden	0		0		0		0		0		0	
Switzerland	1		1		1		0		0		0	
UK	1		1		1		1		1		1	
USA	5	0.95	5	0.87	5	-0.66	5	0.28	5	0.26	5	0.97
USSR	0		0		0		0		0		0	
Yugoslavia	0		0		0		0		0		0	
UK (Manufactured Cigs)	5	0.96	5	0.90	5	0.89	5	-0.73	5	0.90	5	0.52

Table R5.37
Summary of Pearson correlation coefficients, relating Respiratory Disease mortality to prevalence of smoking 20 years earlier, within 30 countries.

All tobacco products

	Males						Females					
	45-49		60-64		75-79		45-49		60-64		75-79	
	N	R	N	R	N	R	N	R	N	R	N	R
Australia	1		1		1		1		1		1	
Austria	5	0.82	5	0.55	5	0.84	5	-0.27	5	0.87	5	-0.21
Belgium	1		1		1		1		1		1	
Bulgaria	0		0		0		0		0		0	
Canada	4	1.00	4	0.99	4	0.56	2		2		2	
Czechoslovakia	0		0		1		0		0		0	
Denmark	5	0.37	5	0.04	5	0.95	5	0.32	5	0.99	5	0.97
Finland	2		2		3	0.95	1		1		1	
France	5	0.95	5	1.00	5	0.98	5	-0.92	5	-0.79	5	-0.95
Germany	6	0.97	6	0.93	6	0.99	6	-0.86	6	0.44	6	-0.12
Greece	4	0.46	4	0.89	4	0.83	2		2		2	
Hungary	2		2		3	1.00	2		2		2	
Iceland	0		0		0		0		0		0	
Ireland	3	0.92	2		2		2		2		2	
Israel	3	-0.42	3	0.97	3	0.57	2		2		2	
Italy	6	0.99	6	0.98	6	0.93	6	-0.95	6	-0.98	6	-0.87
Japan	6	0.96	6	0.66	6	-0.30	6	-0.51	6	0.87	6	0.36
Netherlands	1		1		2		1		1		1	
NewZealand	3	0.89	3	1.00	3	0.94	3	0.97	3	0.94	3	-0.89
Norway	5	0.70	5	0.33	5	0.15	5	-0.03	5	1.00	5	0.98
Poland	4	0.81	4	0.84	4	0.98	4	-0.97	4	-0.94	4	-0.96
Portugal	1		1		1		1		1		1	
Romania	2		2		2		2		2		2	
Spain	2		2		2		2		2		2	
Sweden	3	1.00	3	0.66	3	-1.00	3	0.98	3	0.80	3	1.00
Switzerland	3	0.79	3	-1.00	3	0.55	0		0		0	
UK	11	0.97	11	0.95	11	0.30	5	-0.52	5	0.60	5	0.89
USA	9	0.78	9	0.20	9	-0.11	9	-0.06	9	0.58	9	0.96
USSR	0		0		0		0		0		0	
Yugoslavia	1		1		2		0		0		0	